

ALGEBRA QUALIFYING EXAMINATION

JANUARY 2014

Do either one of nA or nB for $1 \leq n \leq 5$. Justify all your answers. Say what you mean, mean what you say.

- 1A. Let A be an $n \times n$ real skew-symmetric matrix.
- Prove that the nonzero eigenvalues of A are purely imaginary.
 - Prove that $\det(A + I_n) \geq 1$.
- 1B. Let F be a field, and let $M_n(F)$ denote the ring of n -by- n matrices in F . Let $A \in M_n(F)$ with minimal polynomial equal to its characteristic polynomial. Suppose that $B \in M_n(F)$ commutes with A . Show that $B = f(A)$ for some $f \in F[x]$.
- 2A. Prove that a finite simple group of even order is generated by elements of order 2.
- 2B. Show that all groups of order $5 \cdot 7 \cdot 73$ are cyclic.
- 3B. Consider the ring $R = \mathbb{Z}[x]/(x^2)$.
- Show that every ideal of R can be generated by two or fewer elements.
 - Show that (x) is the only prime ideal of R that is not maximal.
- 4A. Let $p_1 < p_2 < \dots < p_r$ be positive prime numbers for some $r \geq 1$, and let K be the field extension of \mathbb{Q} obtained by adjoining $\sqrt{p_i}$ for $1 \leq i \leq r$.
- Prove that K is a Galois extension of \mathbb{Q} with $\text{Gal}(K/\mathbb{Q})$ an elementary abelian 2-group.
 - If $E \subset K$ is a subfield of K of degree 2 over \mathbb{Q} , prove that $E = \mathbb{Q}(\sqrt{m})$ for some m that is the product of all elements in a subset of $\{p_1, p_2, \dots, p_r\}$.
- 4B. Let p be a prime, and let F be a field of characteristic not equal to p . Suppose that F contains a nontrivial p th root of unity, and let $a \in F^\times$.
- Show that a is a p th power in F^\times if and only if $x^p - a$ is reducible in $F[x]$.
 - Let E be the splitting field of $x^p - a$, and suppose that a is not a p th power in F^\times . Determine $\text{Gal}(E/F)$ up to isomorphism.
- 5A. Let R be a ring with unity. Let M be a left R -module in which the chains of distinct left R -submodules are of finite and bounded length. Let $f: M \rightarrow M$ be a left R -module endomorphism. Show that there exist f -stable left R -submodules U and N of M with $M = N \oplus U$ such that f restricts to an isomorphism of U and f restricts to a nilpotent endomorphism of N (i.e., $f^k(N) = 0$ for k sufficiently large).
- 5B. Let $f \in \mathbb{Q}[x]$ be a polynomial of degree $n \geq 1$ with exactly r real roots in \mathbb{C} .
- Show that f factors in $\mathbb{R}[x]$ as a product of r linear and $(n - r)/2$ quadratic polynomials. (You may use the fundamental theorem of algebra.)
 - Let $K = \mathbb{Q}[x]/(f)$. Show that

$$\mathbb{R} \otimes_{\mathbb{Q}} K \cong \mathbb{R}^r \times \mathbb{C}^{(n-r)/2}.$$